



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

THE TWO RACES OF *SAXICOLA ÆNANTHE*.

BY LEONHARD STEJNEGER.

WHEN preparing my paper 'On the Wheatears (*Saxicola*) occurring in North America' (Proc. U. S. Nat. Mus., XXIII, No. 1220, p. 473-481) in which I have demonstrated the existence in this hemisphere of two forms, viz., *Saxicola ænanthe* in Alaska and *Saxicola ænanthe leucorhoa* in Greenland, I wrote to my friend Dr. R. Bowdler Sharpe for the wing measurement of certain African specimens in the British Museum. The answer came too late to be incorporated in the paper quoted, but the facts contained in it are so interesting that I take this means of publishing them and thus further strengthen the claims of the two races for recognition.

The measurements kindly sent by Dr. Sharpe, for which I hereby extend my cordial thanks, are as follows:

*Saxicola ænanthe.*

1 ♂	Ashi Kisa, Interior of British East Africa. Lord Delamere coll. . . . .	Wing 94 mm.
6 ♂♂	Ashi Kisa, Interior of British East Africa. Lord Delamere coll. . . . .	" 97-99 "
1 ♀	Ashi Kisa, Interior of British East Africa. Lord Delamere coll. . . . .	" 94 "
5 ♂♂	British East Africa, F. J. Jackson coll. . . . .	" 95-97 "
2 ♀♀	British East Africa, F. J. Jackson coll. . . . .	" 90-97 "
Total 16	specimens from East Africa . . . . .	" 90-99 "
Average	. . . . .	" 94.5 "

*Saxicola ænanthe leucorhoa.*

1 ad.	Gambia. West Africa . . . . .	Wing 104. mm.
-------	-------------------------------	---------------

These figures, it will be seen, fully substantiate my contention that there are two races, one with the wing normally not exceeding 100 mm., and one with the wing normally not shorter than 100 mm., and that the latter is confined to the extreme west of Europe

and Africa migrating in summer to Greenland and adjacent countries to breed. In the paper quoted I stated "that out of a total of 122 typical *Saxicola ænanthe* only 5 have the wing 100 or 101 mm., while of 45 *Saxicola leucorhoa* none measure less than 100 mm. In other words, only 4 per cent of the small race exceed 99 mm., while none of the larger are below 100. In the whole series of 165 birds, consequently, only 3 per cent of the specimens are intermediate." This statement can now be modified to the effect that of a total of 138 typical *S. ænanthe* only 3.6 per cent of the small race exceed 99 mm., while of 46 specimens of the larger race none are below 100 mm., and that consequently out of the whole series of 182 birds only 2.75 per cent are intermediate.

It is also interesting to note that the average of the extremes of Dr. Sharpe's measurements of the small race, viz., 94.5 mm., is identical with the average of the 65 specimens measured by me as shown in the diagram (*op. cit.*, p. 481) and that the West African specimen of the larger race is identical with the average of 28 specimens as shown in the same diagram. There is consequently every reason to conclude that the results deducible from the diagram in question are founded in Nature.

It is then pretty safe to say, that the short-winged *Saxicola ænanthe* in Alaska goes to India in winter, and that the long-winged *S. ænanthe leucorhoa* from Greenland winters in West Africa.